

organic compounds

Acta Crystallographica Section C
Crystal Structure
Communications

ISSN 0108-2701

S—H···S hydrogen-bond chain in
thiosalicylic acid


Thomas Steiner

Institut für Chemie – Kristallographie, Freie Universität Berlin, Takustrasse 6,
D-14195 Berlin, Germany
Correspondence e-mail: steiner@chemie.fu-berlin.deReceived 25 January 2000
Accepted 6 April 2000

In crystalline thiosalicylic acid (2-mercaptobenzoic acid), $C_7H_6O_2S$, the carboxylic acid groups form hydrogen-bonded dimers, whereas the S—H groups form an infinite S—H···S—H···S—H hydrogen-bond chain, with an S···S distance of 3.986 (3) Å.

Comment

Although S—H is a classical hydrogen-bonding functional group, there is surprisingly little structural information on S—H···X hydrogen bonds. Even for the most common variant, S—H···O, the published material is scarce (Allen *et al.*, 1997). For all the other cases, like S—H···S, S—H···N, S—H···Cl⁻, S—H···Ph *etc.*, there are only a handful of relevant crystal structures (surveyed by Desiraju & Steiner, 1999; see also Steiner, 1998; Rozenberg *et al.*, 1999). In this context, the crystal structure of thiosalicylic acid, (I), was determined. Initially, it was unclear which kind of hydrogen bond the S—H group would form (inter- or intramolecular; S—H···O or S—H···S).



The molecular structure of (I) in the crystal is shown in Fig. 1. The mercapto and carboxyl groups have normal geometries [C2—S1 = 1.766 (3), S1—H1 = 1.30 (5) Å and C2—S1—H1 = 95 (2)°; O1=C7 = 1.219 (4) and O2—C7 = 1.306 (4) Å]. No intramolecular S—H···O hydrogen bond is formed, but the S—H group is oriented away from the carboxylic acid moiety [torsion angle C1—C2—S1—H1 = 178 (2)°]. This contrasts with the oxygen analogue salicylic acid, where an intramolecular O—H···O=C hydrogen bond is formed with O···O = 2.62 Å (Sundaralingam & Jensen, 1965). The molecule of (I) is significantly non-planar, with the carboxylic acid group rotated 10.0 (2)° out of the aromatic plane. This way, the O1 atom is displaced by 0.131 (5) Å from the ring plane. Atom S1 is displaced from that plane in the opposite direction by -0.016 (4) Å. The angle C1—C2—S1 is




Figure 1
The molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level.

widened to 122.1 (2)°. These distortions indicate significant steric repulsion between O1 and S1 [distance 2.752 (3) Å]. In order to understand why no intramolecular hydrogen bond is formed, one must consider that S···O distances in S—H···O hydrogen bonds have only been found longer than 3.2 Å (Desiraju & Steiner, 1999). A hypothetical S—H···O interaction with an S···O separation of around 2.75 Å would (if relatively linear) have an H···O distance so short that it is deep in the repulsive regime of the hydrogen-bond potential.

The hydrogen-bond scheme that is actually formed in (I) is shown in Fig. 2 and geometrical parameters are listed in Table 1. The carboxylic acid groups form dimers with standard geometry [O···O = 2.657 (4) Å]. The S—H groups form an infinite S—H···S—H···S—H hydrogen-bond chain in the direction of the *y* axis, with distances S···S = 3.986 (3) Å and H···S = 2.72 (5) Å. Since the C—S—H group has refined almost to typical neutron diffraction geometry [*e.g.* S—H = 1.338 (2) Å and C—S—H = 96.9 (2)° in the neutron diffraction




Figure 2
Section of the crystal packing illustrating the system of O—H···O and S—H···S hydrogen bonds. The S—H···S chain runs in the direction of the *y* axis, which is horizontal here. The projection plane is rotated by 10° around the *x* axis from the orientation shown.

study of *N*-acetyl-L-cysteine; Takusagawa *et al.*, 1981], the geometry in Table 1 is already realistic and normalization is unnecessary. The distances are similar to those in other S—H...S hydrogen bonds between poorly activated S—H groups, like in thiols and thiophenols (they can be considerably shorter between more polar S—H groups). The acceptor directionality is described by the angle H...S1—C2 of 76.1 (9)° and the torsion angle H...S1—C2—C1 of -81 (1)°, showing that the hydrogen bonds are directed at the 'sides' of the C—S—H acceptors. Two weak C—H...O hydrogen bonds are also formed (Table 1).

Experimental

Thiosalicylic acid, (I), was obtained from Aldrich and was recrystallized from MeOH by slow evaporation of the solvent.

Crystal data

$C_7H_6O_2S$	$D_x = 1.489 \text{ Mg m}^{-3}$
$M_r = 154.18$	Mo $K\alpha$ radiation
Monoclinic, $P2_1/c$	Cell parameters from 34 reflections
$a = 7.856 (7) \text{ \AA}$	$\theta = 7.2\text{--}16.2^\circ$
$b = 5.963 (5) \text{ \AA}$	$\mu = 0.396 \text{ mm}^{-1}$
$c = 14.940 (11) \text{ \AA}$	$T = 293 (2) \text{ K}$
$\beta = 100.69 (7)^\circ$	Plate, yellow
$V = 687.7 (10) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.03 \text{ mm}$
$Z = 4$	

Data collection

Stoe four-circle diffractometer	$h = -10 \rightarrow 7$
ω scans	$k = -7 \rightarrow 7$
1692 measured reflections	$l = -19 \rightarrow 14$
1580 independent reflections	3 standard reflections
1188 reflections with $I > 2\sigma(I)$	frequency: 90 min
$R_{\text{int}} = 0.043$	intensity decay: 5%
$\theta_{\text{max}} = 27.50^\circ$	

Table 1

Hydrogen-bonding geometry (\AA , $^\circ$).

$D\text{--}H\cdots A$	$D\text{--}H$	$H\cdots A$	$D\cdots A$	$D\text{--}H\cdots A$
O2—H2...O1 ⁱ	0.91 (6)	1.75 (6)	2.657 (4)	174 (5)
S1—H1...S1 ⁱⁱ	1.30 (5)	2.72 (5)	3.986 (3)	164 (3)
C5—H5...O1 ⁱⁱⁱ	0.93	2.66	3.584 (5)	171
C6—H6...O2 ^{iv}	0.93	2.62	3.362 (5)	137

Symmetry codes: (i) $1-x, -1-y, -z$; (ii) $1-x, \frac{1}{2}+y, \frac{1}{2}-z$; (iii) $x-1, y, z$; (iv) $-x, -1-y, -z$.

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.171$
 $S = 1.026$
 1580 reflections
 103 parameters
 H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.1042P)^2 + 0.1754P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.53 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

H atoms bonded to C atoms were treated in the default riding model; the H atoms bonded to O2 and S1 were located in difference Fourier calculations and refined isotropically. All H-atom displacement parameters refined to realistic values [H atoms bonded to C: 0.056–0.072 \AA^2 ; H atoms bonded to O and S: 0.11 (1) and 0.09 (1) \AA^2 , respectively].

Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1985); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 1990) and *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

Dr D. Lentz is thanked for providing access to the single-crystal diffractometer.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: DE1138). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H., Bird, C. M., Rowland, R. S. & Raithby, P. R. (1997). *Acta Cryst.* B53, 696–701.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology*, pp. 253–266. Oxford University Press.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Rozenberg, M. S., Nishio, T. & Steiner, T. (1999). *New J. Chem.* 23, 585–586.
- Sheldrick, G. M. (1985). *SHELXS86*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (1990). *Acta Cryst.* A45, C-34.
- Steiner, T. (1998). *Acta Cryst.* B54, 456–463.
- Sundaralingam, M. & Jensen, L. H. (1965). *Acta Cryst.* 18, 1053–1058.
- Takusagawa, F., Koetzle, T. F., Kou, W. W. H. & Parthasarathy, R. (1981). *Acta Cryst.* B37, 1591–1596.

```

#####
###                                     ###
###           Full Paper (Acta Crystallographica Section C)           ###
###                                     ###
#####
#
# This CIF contains the data in a paper accepted for publication in Acta
# Crystallographica Section C. It conforms to the editorial and technical
# requirements of Notes for Authors for Section C, and has been peer
# reviewed under the auspices of the IUCr Commission on Journals.
#
# Full details of the Crystallographic Information File format
# are given in the paper "The Crystallographic Information File (CIF):
# a New Standard Archive File for Crystallography" by S. R. Hall, F. H.
# Allen and I. D. Brown [Acta Cryst. (1991), A47, 655-685].
#
# The current version of the core CIF dictionary is obtainable from
# ftp://ftp.iucr.org/pub/cif_core.dic. The current version number is 2.0.1.
#
# Software is freely available for graphical display of the structure(s) in
# this CIF. For information consult the CIF home page http://www.iucr.org/
# cif/home.html
#
#           Copyright International Union of Crystallography
#
#####
data_salic
_audit_creation_method      SHELXL-97
_journal_date_recd_electronic 2000-01-25
_journal_date_accepted      2000-04-06
_journal_name_full          'Acta Crystallographica, Section C'
_journal_year               2000
_journal_volume             56
_journal_issue              7
_journal_page_first         876
_journal_page_last         877
_journal_paper_category     FO
_publ_contact_author_name   'Dr. Thomas Steiner'
_publ_contact_author_address
;
Institut fur Chemie - Kristallographie
Freie Universitat Berlin
Takustrasse 6
D-14195 Berlin
Germany
;
_publ_contact_author_email   steiner@chemie.fu-berlin.de
_publ_section_title
;
S-H...S hydrogen-bond chain in thiosalicylic acid
;
loop_
  _publ_author_name
  _publ_author_address
  'Steiner, Thomas'
;
Institut f\"ur Chemie - Kristallographie
Freie Universit\"at Berlin
Takustrasse 6
D-14195 Berlin
Germany
;
data_I
_chemical_name_systematic
;
2-mercaptobenzoic acid
;
_chemical_name_common       'thiosalicylic acid'
_chemical_formula_moiety    'C7 H6 O2 S'
_chemical_formula_sum       'C7 H6 O2 S'
_chemical_formula_weight    154.18
_chemical_melting_point     ?
_symmetry_cell_setting      monoclinic
_symmetry_space_group_name_H-M 'P 21/c'
loop_
  _symmetry_equiv_pos_as_xyz
  'x, y, z'
  '-x, y+1/2, -z+1/2'
  '-x, -y, -z'
  'x, -y-1/2, z-1/2'
_cell_length_a              7.856(7)
_cell_length_b              5.963(5)
_cell_length_c              14.940(11)
_cell_angle_alpha           90.00
_cell_angle_beta            100.69(7)

```

```

_cell_angle_gamma      90.00
_cell_volume          687.7(10)
_cell_formula_units_Z  4
_cell_measurement_reflns_used    34
_cell_measurement_theta_min      7.2
_cell_measurement_theta_max     16.2
_cell_measurement_temperature    293(2)
_exptl_crystal_description      plate
_exptl_crystal_colour           yellow
_exptl_crystal_size_max         .25
_exptl_crystal_size_mid         .20
_exptl_crystal_size_min         .03
_exptl_crystal_density_diffn    1.489
_exptl_crystal_density_meas     ?
_exptl_crystal_density_method   'not measured'
_exptl_crystal_F_000           320
_exptl_absorpt_coefficient_mu   .396
_exptl_absorpt_correction_type  none
_exptl_absorpt_process_details  ?
_exptl_absorpt_correction_T_min ?
_exptl_absorpt_correction_T_max ?
_exptl_special_details
;
?
;
_diffn_ambient_temperature      293(2)
_diffn_radiation_type           MoK\alpha
_diffn_radiation_wavelength     .71073
_diffn_radiation_source         'fine-focus sealed tube'
_diffn_radiation_monochromator   graphite
_diffn_measurement_device_type   'Stoe four circle diffractometer'
_diffn_measurement_method       '\w-scans'
_diffn_detector_area_resol_mean  ?
_diffn_reflns_number            1692
_diffn_reflns_av_R_equivalents  .0432
_diffn_reflns_av_sigmaI/netI    .0458
_diffn_reflns_theta_min         2.64
_diffn_reflns_theta_max         27.50
_diffn_reflns_theta_full        27.50
_diffn_measured_fraction_theta_max 1.000
_diffn_measured_fraction_theta_full 1.000
_diffn_reflns_limit_h_min       -10
_diffn_reflns_limit_h_max        7
_diffn_reflns_limit_k_min        -7
_diffn_reflns_limit_k_max        7
_diffn_reflns_limit_l_min       -19
_diffn_reflns_limit_l_max        14
_diffn_standards_number          3
_diffn_standards_interval_count  ?
_diffn_standards_interval_time   90
_diffn_standards_decay_%         5
_refine_special_details
;

```

Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

```

;
_reflns_number_total            1580
_reflns_number_gt               1188
_reflns_threshold_expression     I>2\s(I)
_refine_ls_structure_factor_coef Fsqd
_refine_ls_matrix_type          full
_refine_ls_R_factor_all          .0810
_refine_ls_R_factor_gt           .0567
_refine_ls_wR_factor_ref         .1710
_refine_ls_goodness_of_fit_ref   1.026
_refine_ls_restrained_S_all      1.026
_refine_ls_number_reflns         1580
_refine_ls_number_parameters     103
_refine_ls_number_restraints     0
_refine_ls_hydrogen_treatment    mixed
_refine_ls_weighting_scheme      calc
_refine_ls_weighting_details
'calc w = 1/[\s^2^(Fo^2)+(0.1042P)^2+0.1754P] where P=(Fo^2+2Fc^2)/3'
_refine_ls_shift/su_max          <0.001
_refine_ls_shift/su_mean         .000
_refine_diff_density_max         .530
_refine_diff_density_min        -.303
_refine_ls_extinction_method     none
_refine_ls_extinction_coef       ?
loop_
  _atom_type_symbol

```

```

_atom_type_description
_atom_type_scatter_dispersion_real
_atom_type_scatter_dispersion_imag
_atom_type_scatter_source
'C' 'C' .0033 .0016
'H' 'H' .0000 .0000
'O' 'O' .0106 .0060
'S' 'S' .1246 .1234
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
_computing_data_collection ?
_computing_cell_refinement ?
_computing_data_reduction ?
_computing_structure_solution 'SHELXS86 (Sheldrick, 1986)'
_computing_structure_refinement 'SHELXL97 (Sheldrick, 1997)'
_computing_molecular_graphics 'PLATON (Spek, 1990)'
_computing_publication_material 'SHELXL97'

```

```

loop_
_atom_site_label
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_calc_flag
_atom_site_refinement_flags
_atom_site_occupancy
_atom_site_disorder_assembly
_atom_site_disorder_group
_atom_site_type_symbol
S1 .51689(9) .11696(13) .16353(5) .0462(3) Uani d . 1 . . S
H1 .495(6) .301(8) .206(3) .089(14) Uiso d . 1 . . H
O1 .5120(3) -.2870(4) .07396(16) .0513(6) Uani d . 1 . . O
O2 .2759(3) -.4588(4) .00178(18) .0567(7) Uani d . 1 . . O
H2 .355(7) -.542(11) -.021(3) .108(18) Uiso d . 1 . . H
C1 .2377(3) -.1588(5) .09739(18) .0364(6) Uani d . 1 . . C
C2 .2970(3) .0356(5) .14519(18) .0368(6) Uani d . 1 . . C
C3 .1783(4) .1685(6) .1799(2) .0495(8) Uani d . 1 . . C
H3 .2164 .2970 .2129 .072(12) Uiso calc R 1 . . H
C4 .0063(4) .1121(6) .1661(2) .0516(8) Uani d . 1 . . C
H4 -.0712 .2041 .1889 .056(10) Uiso calc R 1 . . H
C5 -.0528(4) -.0779(6) .1191(2) .0501(8) Uani d . 1 . . C
H5 -.1694 -.1161 .1106 .065(11) Uiso calc R 1 . . H
C6 .0623(4) -.2121(6) .0845(2) .0437(7) Uani d . 1 . . C
H6 .0221 -.3405 .0520 .057(10) Uiso calc R 1 . . H
C7 .3548(4) -.3061(5) .05711(19) .0400(6) Uani d . 1 . . C

```

```

loop_
_atom_site_aniso_label
_atom_site_aniso_U_11
_atom_site_aniso_U_22
_atom_site_aniso_U_33
_atom_site_aniso_U_12
_atom_site_aniso_U_13
_atom_site_aniso_U_23
S1 .0355(4) .0408(5) .0627(5) -.0039(3) .0104(3) -.0065(3)
O1 .0319(10) .0506(13) .0726(14) -.0017(9) .0126(9) -.0245(11)
O2 .0367(11) .0542(14) .0810(17) -.0036(10) .0160(11) -.0309(12)
C1 .0335(13) .0366(14) .0406(13) .0022(11) .0104(10) -.0005(11)
C2 .0323(13) .0387(14) .0401(13) .0018(11) .0088(10) .0001(11)
C3 .0440(16) .0448(17) .0615(19) .0036(13) .0147(14) -.0144(14)
C4 .0418(16) .056(2) .0611(19) .0086(14) .0199(14) -.0111(15)
C5 .0329(14) .063(2) .0569(18) -.0003(13) .0139(12) -.0068(15)
C6 .0347(14) .0460(16) .0515(16) -.0029(12) .0114(11) -.0088(13)
C7 .0369(14) .0369(14) .0478(15) -.0007(12) .0122(11) -.0051(12)

```

```

;geom_special_details
;

```

All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

```

;
loop_
_geom_bond_atom_site_label_1
_geom_bond_atom_site_label_2
_geom_bond_site_symmetry_2
_geom_bond_distance
_geom_bond_publ_flag
S1 C2 . 1.766(3) ?
S1 H1 . 1.30(5) ?
O1 C7 . 1.219(4) ?
O2 C7 . 1.306(4) ?
O2 H2 . .91(6) ?

```

C1 C6 . 1.392(4) ?
C1 C2 . 1.395(4) ?
C1 C7 . 1.478(4) ?
C2 C3 . 1.396(4) ?
C3 C4 . 1.370(4) ?
C3 H3 . .9300 ?
C4 C5 . 1.368(5) ?
C4 H4 . .9300 ?
C5 C6 . 1.377(4) ?
C5 H5 . .9300 ?
C6 H6 . .9300 ?

loop_

_geom_angle_atom_site_label_1
_geom_angle_atom_site_label_2
_geom_angle_atom_site_label_3
_geom_angle_site_symmetry_1
_geom_angle_site_symmetry_3
_geom_angle
_geom_angle_publ_flag
C2 S1 H1 . . 95(2) ?
C7 O2 H2 . . 110(3) ?
C6 C1 C2 . . 119.2(3) ?
C6 C1 C7 . . 118.9(3) ?
C2 C1 C7 . . 121.9(2) ?
C1 C2 C3 . . 118.6(3) ?
C1 C2 S1 . . 122.1(2) ?
C3 C2 S1 . . 119.2(2) ?
C4 C3 C2 . . 120.9(3) ?
C4 C3 H3 . . 119.6 ?
C2 C3 H3 . . 119.6 ?
C5 C4 C3 . . 120.8(3) ?
C5 C4 H4 . . 119.6 ?
C3 C4 H4 . . 119.6 ?
C4 C5 C6 . . 119.3(3) ?
C4 C5 H5 . . 120.3 ?
C6 C5 H5 . . 120.3 ?
C5 C6 C1 . . 121.2(3) ?
C5 C6 H6 . . 119.4 ?
C1 C6 H6 . . 119.4 ?
O1 C7 O2 . . 122.7(3) ?
O1 C7 C1 . . 122.9(3) ?
O2 C7 C1 . . 114.4(3) ?

loop_

_geom_torsion_atom_site_label_1
_geom_torsion_atom_site_label_2
_geom_torsion_atom_site_label_3
_geom_torsion_atom_site_label_4
_geom_torsion_site_symmetry_1
_geom_torsion_site_symmetry_2
_geom_torsion_site_symmetry_3
_geom_torsion_site_symmetry_4
_geom_torsion
_geom_torsion_publ_flag
C6 C1 C2 C3 1.1(4) ?
C7 C1 C2 C3 178.9(3) ?
C6 C1 C2 S1 -179.3(2) ?
C7 C1 C2 S1 -1.5(4) ?
C1 C2 C3 C4 -1.2(5) ?
S1 C2 C3 C4 179.2(3) ?
C2 C3 C4 C5 1.1(5) ?
C3 C4 C5 C6 -.8(5) ?
C4 C5 C6 C17(5) ?
C2 C1 C6 C5 -.8(5) ?
C7 C1 C6 C5 -178.7(3) ?
C6 C1 C7 O1 -171.1(3) ?
C2 C1 C7 O1 11.1(5) ?
C6 C1 C7 O2 9.0(4) ?
C2 C1 C7 O2 -168.8(3) ?

loop_

_geom_hbond_atom_site_label_D
_geom_hbond_atom_site_label_H
_geom_hbond_atom_site_label_A
_geom_hbond_site_symmetry_A
_geom_hbond_distance_DH
_geom_hbond_distance_HA
_geom_hbond_distance_DA
_geom_hbond_angle_DHA
O2 H2 O1 3_645 .91(6) 1.75(6) 2.657(4) 174(5)
S1 H1 S1 2_655 1.30(5) 2.72(5) 3.986(3) 164(3)
C5 H5 O1 1_455 .93 2.66 3.584(5) 171
C6 H6 O2 3_545 .93 2.62 3.362(5) 137